



Bowel Management

Policies & Procedures

Shop 5, 38 Princess Street
Bundaberg East Qld 4670

07 4361 6848

www.123supports.com

www.facebook.com/123supports

ABN: 14 930 943 229

TABLE OF CONTENT

Policy / Procedure	Page
BOWEL MANAGEMENT	3
Scope	3
Purpose.....	3
Constipation	3
Diarrhoea	4
Faecal incontinence	4
Seek medical attention.....	5
Promote good bowel management.....	5
Bowel care guidelines	5
Further Advice or Assistance	5
Effectiveness and Review.....	5

BOWEL MANAGEMENT

Scope

When

- Applies when clients require ongoing bowel management

Who

- Applies to all employees, supervisors and key management personnel supporting clients with ongoing bowel management needs

Purpose

This policy sets the principles for providing bowel care to people we support. Quality bowel care is about ensuring a person's bowels are effectively managed in a safe, consistent, and hygienic manner that promotes a person's overall health and well-being. All clients we support deserve:

- Appropriate advice
- Monitoring
- Support and intervention

Above all, clients must be treated with dignity at all times regarding a sensitive and important bodily function.

A healthy bowel can be affected by a number of factors including:

- A diet lacking in fibre
- Insufficient fluid intake
- Disruption to regular diet or routine
- Delaying bowel actions due to pain (such as haemorrhoids)
- Certain types of medications – narcotics, opioids, sedatives, codeine-based medications will slow down bowel movements
- Recent illness or hospitalisation
- Low activity levels and reduced mobility – particularly those who are in wheelchairs, bed-bound or post-surgery
- Reduced physical and emotional wellbeing

People with disability are more vulnerable to suffer from bowel dysfunction (a bowel that is not functioning correctly) for any of the reasons described above. Bowel function can also be affected by diabetes, obesity, heart problems and other conditions. The main types of bowel dysfunction are:

- Constipation and poor bowel emptying
- Diarrhoea
- Faecal incontinence

Constipation

Constipation is a condition in which bowel movements occur less often than usual or consist of hard, dry stools that are painful or difficult to pass. While bowel habits vary, an adult who has not had a bowel movement in three days or a child who has not had a bowel movement in four days is considered constipated.

Constipation has many possible causes, including low mobility, inadequate diet, slow movement of faeces through bowels, or abnormality of the muscles involved in emptying the bowel.

Preventing constipation is best practice and involves ensuring a person:

- Has sufficient fibre in their diet – see list of high residue foods
- Relies less on foods low in fibre (low residue foods)
- Has adequate intake of fluids (typically 6-8 glasses of water a day)
- Reduces or avoids intake of caffeine drinks (tea, coffee, and cola)
- Has regular exercise
- Goes to the toilet when the urge is felt and has enough time

Low residue foods (low fibre foods slow down transit times)	High residue foods (high fibre foods speed up transit times)
<ul style="list-style-type: none">• White bread, white rice, pasta• Raw lettuce, cucumber, avocado• Cooked vegetables without skin or seeds• Clear broths and strained soups• Raw fruit such as apricots, bananas, rock melon, honeydew, watermelon, nectarines, papayas, peaches, plums• Tender-cooked meat, seafood, eggs• Canned vegetables and fruit• Butter, vegetable oils, margarine• Yoghurt, cottage cheese, ricotta cheese, milk, soy milk, tofu• • Fruit juice without pulp	<ul style="list-style-type: none">• Wholegrain cereals, oats, brown rice, bread or cakes with bran, nut or seeds, wholemeal pasta• Most leafy vegetables, broccoli, cauliflower• Raw fruit and vegetables with skins and seeds• Nuts, legumes, corn, popcorn• Dried fruit, prunes• Fruit juice with pulp, prune juice

Diarrhoea

Diarrhoea is defined as more than 300mls of loose watery stools in a 24-hour period. Diarrhoea can be short lived such as with food poisoning or bowel infection (which can be spread to other people), or it can be as a result of:

- Inflammatory bowel disease
- Irritable bowel syndrome
- Coeliac disease
- Crohn's disease

Diarrhoea may also be caused by food allergies, medications, radiation therapy, overuse of laxatives and diabetes. Consult a doctor if a client suffers ongoing diarrhoea.

Faecal incontinence

Faecal incontinence is leakage from the bowel due to poor bowel control. Incontinence can often be experienced by people with health conditions such as diabetes, obesity, heart problems, stroke, cancer, prostate and kidney problems, and neurological conditions such as spina bifida, autism, spinal cord injuries and Parkinson's disease.

Incontinence is best managed or prevented with a healthy diet high in fibre, staying active, and regular bowel habits.

Seek medical attention

Call a doctor or seek medical advice if a client experiences any of the following:

- Vomiting blood or faecal matter
- Diarrhoea and/or vomiting that is more than a one-off event
- Bleeding from the bowel
- Fresh (red) or old (black) blood in faeces (note: a person may also have black faeces when taking iron supplements)
- Unusual pain before, during or after bowel action
- Constipation not resolved by medication – bowels not open for three days may indicate:
 - Faecal impaction
 - Bowel obstruction
 - Paralytic ileus

Promote good bowel management

- Ensure clients have sufficient fluid intake
- Ensure clients have sufficient intake of dietary fibre
- Ensure clients have adequate physical activity
- Provide a relaxed and unhurried setting with which clients can move their bowels
- If necessary, provide planning and daily bowel diary for clients with bowel management needs

Bowel care guidelines

- Maintain good hand hygiene
- Use disposable incontinence pads and pants if needed
- Suggest clients wear underwear made of natural fibres such as cotton
- Clean and wash surrounding peri anal skin/rectal region/moist areas of buttocks and groin with warm water and mild unscented soap
- Monitor for skin reaction inflammation, redness, breakdown, bleeding, or erythema

Further Advice or Assistance

Further advice and information can be obtained from the:

- » Director:
 - by phoning: 07 4361 6848;
 - by emailing: admin@123supports.com
- » NDIS Quality and Safeguards Commission (<https://www.ndiscommission.gov.au>)

Effectiveness and Review

The Director will review this Policy and Procedures document each 12 months on the anniversary of its approval.